

Adaptive Filon methods for the computation of highly oscillatory integrals

Marnix Van Daele, Veerle Ledoux

Department of Applied Mathematics and Computer Science
Ghent University

S. Margherita di Pula, Cagliari, Italy, October 10-14 2011

Oscillatory integrals

$$I[f] = \int_0^h f(x) e^{i\omega g(x)} dx$$

We focus on the particular case

$$I[f] = \int_0^h f(x) e^{i\omega x} dx$$

If the integrand oscillates rapidly, and unless we use a huge number of function evaluations, the classical ν -point Gauss rule is useless.

Oscillatory integrals

$$I[f] = \int_0^h f(x) e^{i\omega g(x)} dx$$

We focus on the particular case

$$I[f] = \int_0^h f(x) e^{i\omega x} dx$$

If the integrand oscillates rapidly, and unless we use a huge number of function evaluations, the classical ν -point Gauss rule is useless.

Oscillatory integrals

$$I[f] = \int_0^h f(x) e^{i\omega g(x)} dx$$

We focus on the particular case

$$I[f] = \int_0^h f(x) e^{i\omega x} dx$$


If the integrand oscillates rapidly, and unless we use a huge number of function evaluations, the classical ν -point Gauss rule is useless.



Gauss rule applied to oscillatory integrands

Example : $f(x) = \exp(x)$ and $h = 1/10$

$$\int_0^h e^x e^{i\omega x} dx = \frac{-1 + e^{h(1+i\omega)}}{1 + i\omega}$$



The absolute error in Gauss-Legendre quadrature for different values of the **characteristic frequency** $\psi = \omega h$.

Asymptotic expansion

$$\begin{aligned} I[f] &= \int_a^b f(x) e^{i\omega x} dx \\ &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) - \frac{1}{i\omega} I[f'] \\ &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) \\ &\quad - \frac{1}{(i\omega)^2} \left(f'(b) e^{i\omega b} - f'(a) e^{i\omega a} \right) + \frac{1}{(i\omega)^2} I[f''] \end{aligned}$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

Asymptotic expansion

$$\begin{aligned}
 I[f] &= \int_a^b f(x) e^{i\omega x} dx \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) - \frac{1}{i\omega} I[f'] \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) \\
 &\quad - \frac{1}{(i\omega)^2} \left(f'(b) e^{i\omega b} - f'(a) e^{i\omega a} \right) + \frac{1}{(i\omega)^2} I[f'']
 \end{aligned}$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

Asymptotic expansion

$$\begin{aligned}
 I[f] &= \int_a^b f(x) e^{i\omega x} dx \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) - \frac{1}{i\omega} I[f'] \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) \\
 &\quad - \frac{1}{(i\omega)^2} \left(f'(b) e^{i\omega b} - f'(a) e^{i\omega a} \right) + \frac{1}{(i\omega)^2} I[f'']
 \end{aligned}$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

Asymptotic expansion

$$\begin{aligned}
 I[f] &= \int_a^b f(x) e^{i\omega x} dx \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) - \frac{1}{i\omega} I[f'] \\
 &= \frac{1}{i\omega} \left(f(b) e^{i\omega b} - f(a) e^{i\omega a} \right) \\
 &\quad - \frac{1}{(i\omega)^2} \left(f'(b) e^{i\omega b} - f'(a) e^{i\omega a} \right) + \frac{1}{(i\omega)^2} I[f'']
 \end{aligned}$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

Asymptotic rules

$$I[f] = \int_a^b f(x) e^{i\omega x} dx$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] = - \sum_{m=0}^{s-1} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] - I[f] \sim O(\omega^{-s-1}) \quad \omega \rightarrow +\infty$$

This asymptotic method is of **asymptotic order** $s + 1$.

The asymptotic order gives us the rate at which the error decreases with increasing ω .

Asymptotic rules

$$I[f] = \int_a^b f(x) e^{i\omega x} dx$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] = - \sum_{m=0}^{s-1} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] - I[f] \sim O(\omega^{-s-1}) \quad \omega \rightarrow +\infty$$

This asymptotic method is of **asymptotic order** $s + 1$.

The asymptotic order gives us the rate at which the error decreases with increasing ω .

Asymptotic rules

$$I[f] = \int_a^b f(x) e^{i\omega x} dx$$

$$I[f] = - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] = - \sum_{m=0}^{s-1} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} f^{(m)}(b) - e^{i\omega a} f^{(m)}(a) \right]$$

$$Q_s^A[f] - I[f] \sim O(\omega^{-s-1}) \quad \omega \rightarrow +\infty$$

This asymptotic method is of **asymptotic order** $s + 1$.

The asymptotic order gives us the rate at which the error decreases with increasing ω .

Exponential fitting

M. VAN DAELE, G. VANDEN BERGHE AND H. VANDE VYVER,
*Exponentially fitted quadrature rules of Gauss type for
oscillatory integrands*, Appl. Numer. Math., 53 (2005),
pp. 509–526.

How to compute

$$\int_{-1}^1 F(t) dt$$

whereby $F(x)$ has an oscillatory behaviour with frequency μ ?

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\mu} \int_{-1}^1 f(h(t+1)/2) e^{i\mu t} dt \quad \mu = \frac{\omega h}{2}$$

Exponential fitting

M. VAN DAELE, G. VANDEN BERGHE AND H. VANDE VYVER,
*Exponentially fitted quadrature rules of Gauss type for
 oscillatory integrands*, Appl. Numer. Math., 53 (2005),
 pp. 509–526.

How to compute

$$\int_{-1}^1 F(t) dt$$

whereby $F(x)$ has an oscillatory behaviour with frequency μ ?

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\mu} \int_{-1}^1 f(h(t+1)/2) e^{i\mu t} dt \quad \mu = \frac{\omega h}{2}$$

Exponential fitting

$$\mathcal{L}[F; x; h; \mathbf{a}] = \int_{x-h}^{x+h} F(z) dz - h \sum_{k=1}^{\nu} w_k F(x + \hat{c}_k h), \quad \hat{c}_k \in [-1, 1]$$

(put $x = \mathbf{0}$ and $h = \mathbf{1}$ to obtain $\int_{-1}^1 F(t) dt$)

$\mathcal{L}[F; x; h; \mathbf{a}] = \mathbf{0}$ for a reference set of $K + \mathbf{2}(P + \mathbf{1}) + \mathbf{1} = \mathbf{2}\nu$ functions

$$\mathbf{1}, t, t^2, \dots, t^K,$$

$$\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)$$

In this talk we only consider the case $K = -\mathbf{1}$, $P = \nu - \mathbf{1}$.

Exponential fitting

$$\mathcal{L}[F; x; h; \mathbf{a}] = \int_{x-h}^{x+h} F(z) dz - h \sum_{k=1}^{\nu} w_k F(x + \hat{c}_k h), \quad \hat{c}_k \in [-1, 1]$$

(put $x = \mathbf{0}$ and $h = \mathbf{1}$ to obtain $\int_{-1}^1 F(t) dt$)

$\mathcal{L}[F; x; h; \mathbf{a}] = \mathbf{0}$ for a reference set of $K + \mathbf{2}(P + \mathbf{1}) + \mathbf{1} = \mathbf{2}\nu$ functions

$$\mathbf{1}, t, t^2, \dots, t^K,$$

$$\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)$$

In this talk we only consider the case $K = -\mathbf{1}$, $P = \nu - \mathbf{1}$.

Exponential fitting

$$\mathcal{L}[F; x; h; \mathbf{a}] = \int_{x-h}^{x+h} F(z) dz - h \sum_{k=1}^{\nu} w_k F(x + \hat{c}_k h), \quad \hat{c}_k \in [-1, 1]$$

(put $x = \mathbf{0}$ and $h = \mathbf{1}$ to obtain $\int_{-1}^1 F(t) dt$)

$\mathcal{L}[F; x; h; \mathbf{a}] = \mathbf{0}$ for a reference set of $K + \mathbf{2}(P + \mathbf{1}) + \mathbf{1} = \mathbf{2}\nu$ functions

$$\mathbf{1}, t, t^2, \dots, t^K,$$

$$\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)$$

In this talk we only consider the case $K = -\mathbf{1}$, $P = \nu - \mathbf{1}$.

1-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1)$$

$$\int_{-1}^1 \exp(\pm i\mu x) dx - w_1 \exp(\pm \hat{c}_1 \mu) = 0$$

$$w_1 = 2 \sin(\mu)/\mu \qquad \hat{c}_1 = 0$$

$$I[f] = \int_0^h f(x) \exp(i\omega x) dx = \int_0^h F(x) dx$$

$$Q_1^{EF}[F] = \frac{h \sin(\mu)}{\mu} F(h/2) = \frac{e^{ih\omega} - 1}{i\omega} f(h/2) \quad \mu = \omega h/2$$

1-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1)$$

$$\int_{-1}^1 \exp(\pm i\mu x) dx - w_1 \exp(\pm \hat{c}_1 \mu) = 0$$

$$w_1 = 2 \sin(\mu)/\mu \qquad \hat{c}_1 = 0$$

$$I[f] = \int_0^h f(x) \exp(i\omega x) dx = \int_0^h F(x) dx$$

$$Q_1^{EF}[F] = \frac{h \sin(\mu)}{\mu} F(h/2) = \frac{e^{ih\omega} - 1}{i\omega} f(h/2) \quad \mu = \omega h/2$$

1-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1)$$

$$\int_{-1}^1 \exp(\pm i\mu x) dx - w_1 \exp(\pm \hat{c}_1 \mu) = 0$$

$$w_1 = 2 \sin(\mu)/\mu \qquad \hat{c}_1 = 0$$

$$I[f] = \int_0^h f(x) \exp(i\omega x) dx = \int_0^h F(x) dx$$

$$Q_1^{EF}[F] = \frac{h \sin(\mu)}{\mu} F(h/2) = \frac{e^{ih\omega} - 1}{i\omega} f(h/2) \quad \mu = \omega h/2$$

1-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1)$$

$$\int_{-1}^1 \exp(\pm i\mu x) dx - w_1 \exp(\pm \hat{c}_1 \mu) = 0$$

$$w_1 = 2 \sin(\mu)/\mu \qquad \hat{c}_1 = 0$$

$$I[f] = \int_0^h f(x) \exp(i\omega x) dx = \int_0^h F(x) dx$$

$$Q_1^{EF}[F] = \frac{h \sin(\mu)}{\mu} F(h/2) = \frac{e^{ih\omega} - 1}{i\omega} f(h/2) \quad \mu = \omega h/2$$

2-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1) + w_2 F(\hat{c}_2)$$

$$\begin{cases} \int_{-1}^1 \exp(\pm i \mu x) dx - w_1 \exp(\pm i \hat{c}_1 \mu) - w_2 \exp(\pm i \hat{c}_2 \mu) = 0 \\ \int_{-1}^1 x \exp(\pm i \mu x) dx - w_1 \hat{c}_1 \exp(\pm i \hat{c}_1 \mu) - w_2 \hat{c}_2 \exp(\pm i \hat{c}_2 \mu) = 0 \end{cases}$$

Assuming $w_1 = w_2$ and $\hat{c}_1 = -\hat{c}_2$:

$$\Leftrightarrow \begin{cases} w_2 \mu \cos(\mu \hat{c}_2) - \sin(\mu) = 0 \\ w_2 \hat{c}_2 \mu^2 \sin(\mu \hat{c}_2) - \sin(\mu) + \mu \cos(\mu) = 0 \end{cases}$$

$$Q_2^{EF}[F] = \frac{h}{2} w_2 \left[F\left(\frac{h(1+\hat{c}_2)}{2}\right) + F\left(\frac{h(1-\hat{c}_2)}{2}\right) \right] \quad \mu = \frac{\omega h}{2}$$

2-node EF rule

$$\int_{-1}^1 F(x) dx \approx w_1 F(\hat{c}_1) + w_2 F(\hat{c}_2)$$

$$\begin{cases} \int_{-1}^1 \exp(\pm i \mu x) dx - w_1 \exp(\pm i \hat{c}_1 \mu) - w_2 \exp(\pm i \hat{c}_2 \mu) = 0 \\ \int_{-1}^1 x \exp(\pm i \mu x) dx - w_1 \hat{c}_1 \exp(\pm i \hat{c}_1 \mu) - w_2 \hat{c}_2 \exp(\pm i \hat{c}_2 \mu) = 0 \end{cases}$$

Assuming $w_1 = w_2$ and $\hat{c}_1 = -\hat{c}_2$:

$$\Leftrightarrow \begin{cases} w_2 \mu \cos(\mu \hat{c}_2) - \sin(\mu) = 0 \\ w_2 \hat{c}_2 \mu^2 \sin(\mu \hat{c}_2) - \sin(\mu) + \mu \cos(\mu) = 0 \end{cases}$$

$$Q_2^{EF}[F] = \frac{h}{2} w_2 \left[F\left(\frac{h(1+\hat{c}_2)}{2}\right) + F\left(\frac{h(1-\hat{c}_2)}{2}\right) \right] \quad \mu = \frac{\omega h}{2}$$

2-node EF rule

$$\begin{cases} w_2 \mu \cos(\mu \hat{c}_2) - \sin(\mu) = 0 \\ w_2 \hat{c}_2 \mu^2 \sin(\mu \hat{c}_2) - \sin(\mu) + \mu \cos(\mu) = 0 \end{cases}$$

If $\cos(\mu \hat{c}_2) \neq 0$ then $w_2 = \sin \mu / (\mu \cos(\mu \hat{c}_2))$

$$G(\hat{c}_2) := (\sin \mu - \mu \cos \mu) \cos(\mu \hat{c}_2) - \mu \hat{c}_2 \sin \mu \sin(\mu \hat{c}_2) = 0$$






Figure: $G(x_2)$ for $\mu = 5$, $\mu = 50$ and $\mu = 200$.



2-node EF rule

$$\begin{cases} w_2 \mu \cos(\mu \hat{c}_2) - \sin(\mu) = 0 \\ w_2 \hat{c}_2 \mu^2 \sin(\mu \hat{c}_2) - \sin(\mu) + \mu \cos(\mu) = 0 \end{cases}$$

If $\cos(\mu \hat{c}_2) \neq 0$ then $w_2 = \sin \mu / (\mu \cos(\mu \hat{c}_2))$

$$G(\hat{c}_2) := (\sin \mu - \mu \cos \mu) \cos(\mu \hat{c}_2) - \mu \hat{c}_2 \sin \mu \sin(\mu \hat{c}_2) = 0$$






Figure: $G(x_2)$ for $\mu = 5$, $\mu = 50$ and $\mu = 200$.

2-node EF rule




Figure: The $\hat{c}_2(\mu)$ and $w_2(\mu)$ curve for the EF method with $\nu = 2$.

3-node EF rule

$$\hat{c}_1 = -\hat{c}_3 \quad \hat{c}_2 = 0 \quad w_1 = w_3$$




Figure: The $\hat{c}_3(\mu)$, $w_1(\mu) = w_3(\mu)$ and $w_2(\mu)$ curves for the $\nu = 3$ EF rule

4-node EF rule

$$\hat{c}_1 = -\hat{c}_4 \quad \hat{c}_2 = -\hat{c}_3 \quad w_1 = w_4 \quad w_2 = w_3$$




Figure: Nodes and weights of the EF rule with $\nu = 4$ quadrature nodes.

Accuracy of EF rules

All EF rules reduce to the classical ν -point Gauss(-Legendre) method in the limiting case $\mu = 0$.

Thus for small μ : $O(h^{2\nu+1})$

What about the accuracy for larger values of $\mu = \omega h/2$?

J. P. COLEMAN AND L. GR. IXARU, *Truncation errors in exponential fitting for oscillatory problems*, SIAM. J. Numer. Anal., 44 (2006), pp. 1441–1465.

for large μ : $O(\mu^{\bar{\nu}-\nu})$ with $\bar{\nu} = \lfloor (\nu - 1)/2 \rfloor$

$$\nu = 1 : O(\omega^{-1})$$

$$\nu = 2, 3 : O(\omega^{-2})$$

$$\nu = 4, 5 : O(\omega^{-3})$$

Accuracy of EF rules

All EF rules reduce to the classical ν -point Gauss(-Legendre) method in the limiting case $\mu = 0$.

Thus for small μ : $O(h^{2\nu+1})$

What about the accuracy for larger values of $\mu = \omega h/2$?

J. P. COLEMAN AND L. GR. IXARU, *Truncation errors in exponential fitting for oscillatory problems*, SIAM. J. Numer. Anal., 44 (2006), pp. 1441–1465.

for large μ : $O(\mu^{\bar{\nu}-\nu})$ with $\bar{\nu} = \lfloor (\nu - 1)/2 \rfloor$

$$\nu = 1 : O(\omega^{-1})$$

$$\nu = 2, 3 : O(\omega^{-2})$$

$$\nu = 4, 5 : O(\omega^{-3})$$

Accuracy of EF rules

All EF rules reduce to the classical ν -point Gauss(-Legendre) method in the limiting case $\mu = 0$.

Thus for small μ : $O(h^{2\nu+1})$

What about the accuracy for larger values of $\mu = \omega h/2$?

J. P. COLEMAN AND L. GR. IXARU, *Truncation errors in exponential fitting for oscillatory problems*, SIAM. J. Numer. Anal., 44 (2006), pp. 1441–1465.

for large μ : $O(\mu^{\bar{\nu}-\nu})$ with $\bar{\nu} = \lfloor (\nu - 1)/2 \rfloor$

$$\nu = 1 : O(\omega^{-1})$$

$$\nu = 2, 3 : O(\omega^{-2})$$

$$\nu = 4, 5 : O(\omega^{-3})$$

Accuracy of EF rules

All EF rules reduce to the classical ν -point Gauss(-Legendre) method in the limiting case $\mu = 0$.

Thus for small μ : $O(h^{2\nu+1})$

What about the accuracy for larger values of $\mu = \omega h/2$?

J. P. COLEMAN AND L. GR. IXARU, *Truncation errors in exponential fitting for oscillatory problems*, SIAM. J. Numer. Anal., 44 (2006), pp. 1441–1465.

for large μ : $O(\mu^{\bar{\nu}-\nu})$ with $\bar{\nu} = \lfloor (\nu - 1)/2 \rfloor$

$$\nu = 1 : O(\omega^{-1})$$

$$\nu = 2, 3 : O(\omega^{-2})$$

$$\nu = 4, 5 : O(\omega^{-3})$$

Accuracy of EF rules

All EF rules reduce to the classical ν -point Gauss(-Legendre) method in the limiting case $\mu = 0$.

Thus for small μ : $O(h^{2\nu+1})$

What about the accuracy for larger values of $\mu = \omega h/2$?

J. P. COLEMAN AND L. GR. IXARU, *Truncation errors in exponential fitting for oscillatory problems*, SIAM. J. Numer. Anal., 44 (2006), pp. 1441–1465.

for large μ : $O(\mu^{\bar{\nu}-\nu})$ with $\bar{\nu} = \lfloor (\nu - 1)/2 \rfloor$

$$\nu = 1 : O(\omega^{-1})$$

$$\nu = 2, 3 : O(\omega^{-2})$$

$$\nu = 4, 5 : O(\omega^{-3})$$

Proof

$$\int_{-1}^1 F(t) dt \approx \int_{-1}^1 \bar{F}(t) dt$$

$$\bar{F}(t) \in \text{span}\{\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)\}$$

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\frac{\omega h}{2}} \int_{-1}^1 f\left(\frac{h}{2}(t+1)\right) e^{i\frac{\omega h}{2}t} dt$$

If $\frac{\omega h}{2} = \mu$ then $I[f] \approx I[\bar{f}]$ with $\bar{f}(x) \in \text{span}\{1, x, x^2, \dots, x^{\nu-1}\}$

$$Q_{\nu}^{EF}[f] - I[f] = I[\bar{f}] - I[f] = I[v] \quad v(x) := \bar{f}(x) - f(x)$$

Proof

$$\int_{-1}^1 F(t) dt \approx \int_{-1}^1 \bar{F}(t) dt$$

$$\bar{F}(t) \in \text{span}\{\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)\}$$

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\frac{\omega h}{2}} \int_{-1}^1 f\left(\frac{h}{2}(t+1)\right) e^{i\frac{\omega h}{2}t} dt$$

If $\frac{\omega h}{2} = \mu$ then $I[f] \approx I[\bar{f}]$ with $\bar{f}(x) \in \text{span}\{1, x, x^2, \dots, x^{\nu-1}\}$

$$Q_{\nu}^{EF}[f] - I[f] = I[\bar{f}] - I[f] = I[v] \quad v(x) := \bar{f}(x) - f(x)$$

Proof

$$\int_{-1}^1 F(t) dt \approx \int_{-1}^1 \bar{F}(t) dt$$

$$\bar{F}(t) \in \text{span}\{\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)\}$$

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\frac{\omega h}{2}} \int_{-1}^1 f\left(\frac{h}{2}(t+1)\right) e^{i\frac{\omega h}{2}t} dt$$

$$\text{If } \frac{\omega h}{2} = \mu \text{ then } I[f] \approx I[\bar{f}] \text{ with } \bar{f}(x) \in \text{span}\{1, x, x^2, \dots, x^{\nu-1}\}$$

$$Q_{\nu}^{EF}[f] - I[f] = I[\bar{f}] - I[f] = I[v] \quad v(x) := \bar{f}(x) - f(x)$$

Proof

$$\int_{-1}^1 F(t) dt \approx \int_{-1}^1 \bar{F}(t) dt$$

$$\bar{F}(t) \in \text{span}\{\exp(\pm i\mu t), t \exp(\pm i\mu t), t^2 \exp(\pm i\mu t), \dots, t^P \exp(\pm i\mu t)\}$$

$$I[f] = \int_0^h f(x) e^{i\omega x} dx = \frac{h}{2} e^{i\frac{\omega h}{2}} \int_{-1}^1 f\left(\frac{h}{2}(t+1)\right) e^{i\frac{\omega h}{2}t} dt$$

If $\frac{\omega h}{2} = \mu$ then $I[f] \approx I[\bar{f}]$ with $\bar{f}(x) \in \text{span}\{1, x, x^2, \dots, x^{\nu-1}\}$

$$Q_{\nu}^{EF}[f] - I[f] = I[\bar{f}] - I[f] = I[v] \quad v(x) := \bar{f}(x) - f(x)$$

Proof

Suppose ν is even and $a < c_1 < c_2 < \dots < c_\nu < b$

$$c_j = a + \lambda_j/\omega \quad c_{\nu-j+1} = b - \lambda_j/\omega \quad j = 1, \dots, \nu/2$$

$$v(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{i=1}^{\nu} (x - c_i)$$

$$v(x) = s(x) \prod_{i=1}^{\nu/2} (x - b + \lambda_i/\omega) \quad s(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{j=1}^{\nu/2} (x - a - \lambda_j/\omega)$$

$$v(b) = s(b) \prod_{i=1}^{\nu/2} (\lambda_i/\omega) = O(\omega^{-\nu/2})$$

$$v'(b) = s(b)\omega^{-\nu/2+1} \sum_{k=1}^{\nu/2} \prod_{i \neq k} \lambda_i + O(\omega^{-\nu/2}) = O(\omega^{-\nu/2+1})$$

Proof

Suppose ν is even and $a < c_1 < c_2 < \dots < c_\nu < b$

$$c_j = a + \lambda_j/\omega \quad c_{\nu-j+1} = b - \lambda_j/\omega \quad j = 1, \dots, \nu/2$$

$$v(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{i=1}^{\nu} (x - c_i)$$

$$v(x) = s(x) \prod_{i=1}^{\nu/2} (x - b + \lambda_i/\omega) \quad s(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{j=1}^{\nu/2} (x - a - \lambda_j/\omega)$$

$$v(b) = s(b) \prod_{i=1}^{\nu/2} (\lambda_i/\omega) = O(\omega^{-\nu/2})$$

$$v'(b) = s(b)\omega^{-\nu/2+1} \sum_{k=1}^{\nu/2} \prod_{i \neq k} \lambda_i + O(\omega^{-\nu/2}) = O(\omega^{-\nu/2+1})$$

Proof

Suppose ν is even and $a < c_1 < c_2 < \dots < c_\nu < b$

$$c_j = a + \lambda_j/\omega \quad c_{\nu-j+1} = b - \lambda_j/\omega \quad j = 1, \dots, \nu/2$$

$$v(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{i=1}^{\nu} (x - c_i)$$

$$v(x) = s(x) \prod_{i=1}^{\nu/2} (x - b + \lambda_i/\omega) \quad s(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{j=1}^{\nu/2} (x - a - \lambda_j/\omega)$$

$$v(b) = s(b) \prod_{i=1}^{\nu/2} (\lambda_i/\omega) = O(\omega^{-\nu/2})$$

$$v'(b) = s(b)\omega^{-\nu/2+1} \sum_{k=1}^{\nu/2} \prod_{i \neq k} \lambda_i + O(\omega^{-\nu/2}) = O(\omega^{-\nu/2+1})$$

Proof

Suppose ν is even and $a < c_1 < c_2 < \dots < c_\nu < b$

$$c_j = a + \lambda_j/\omega \quad c_{\nu-j+1} = b - \lambda_j/\omega \quad j = 1, \dots, \nu/2$$

$$v(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{i=1}^{\nu} (x - c_i)$$

$$v(x) = s(x) \prod_{i=1}^{\nu/2} (x - b + \lambda_i/\omega) \quad s(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{j=1}^{\nu/2} (x - a - \lambda_j/\omega)$$

$$v(b) = s(b) \prod_{i=1}^{\nu/2} (\lambda_i/\omega) = O(\omega^{-\nu/2})$$

$$v'(b) = s(b)\omega^{-\nu/2+1} \sum_{k=1}^{\nu/2} \prod_{i \neq k} \lambda_i + O(\omega^{-\nu/2}) = O(\omega^{-\nu/2+1})$$

Proof

Suppose ν is even and $a < c_1 < c_2 < \dots < c_\nu < b$

$$c_j = a + \lambda_j/\omega \quad c_{\nu-j+1} = b - \lambda_j/\omega \quad j = 1, \dots, \nu/2$$

$$v(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{i=1}^{\nu} (x - c_i)$$

$$v(x) = s(x) \prod_{i=1}^{\nu/2} (x - b + \lambda_i/\omega) \quad s(x) = \frac{f^{(\nu)}(\xi(x))}{\nu!} \prod_{j=1}^{\nu/2} (x - a - \lambda_j/\omega)$$

$$v(b) = s(b) \prod_{i=1}^{\nu/2} (\lambda_i/\omega) = O(\omega^{-\nu/2})$$

$$v'(b) = s(b)\omega^{-\nu/2+1} \sum_{k=1}^{\nu/2} \prod_{i \neq k} \lambda_i + O(\omega^{-\nu/2}) = O(\omega^{-\nu/2+1})$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor (\nu-1)/2 \rfloor - \nu}) \end{aligned}$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor(\nu-1)/2\rfloor-\nu}) \end{aligned}$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor(\nu-1)/2\rfloor-\nu}) \end{aligned}$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor(\nu-1)/2\rfloor-\nu}) \end{aligned}$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor (\nu-1)/2 \rfloor - \nu}) \end{aligned}$$

Proof

$$v(b) = O(\omega^{-\nu/2}) \qquad v'(b) = O(\omega^{-\nu/2+1})$$

$$v^{(n)}(b) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$v^{(n)}(a) = O(\omega^{-\nu/2+n}), \quad n = 0, 1, \dots, \nu/2 - 1$$

$$\begin{aligned} Q_\nu^{EF}[f] - I[f] &= I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega b} v^{(m)}(b) - e^{i\omega a} v^{(m)}(a) \right] \\ &= - \sum_{m=0}^{\nu/2-1} \frac{1}{(-i\omega)^{m+1}} O(\omega^{-\nu/2+m}) + O(\omega^{-\nu/2-1}) \\ &= O(\omega^{-\nu/2-1}) = O(\omega^{\lfloor(\nu-1)/2\rfloor-\nu}) \end{aligned}$$

Filon-type

L. N. G. FILON, *On a quadrature formula for trigonometric integrals*, Proc. Royal Soc. Edinburgh, 49 (1928), pp. 38–47.

Interpolate only the function $f(x)$ at $c_1 h, \dots, c_\nu h$ by a polynomial $\bar{f}(x)$

$$I[f] \approx Q_\nu^F[f] = \int_0^h \bar{f}(x) e^{i\omega x} dx = h \sum_{l=1}^{\nu} b_l(ih\omega) f(c_l h)$$

$$b_l(ih\omega) = \int_0^1 \ell_l(x) e^{ih\omega x} dx$$

ℓ_l is the l th cardinal polynomial of Lagrangian interpolation.

Filon-type

L. N. G. FILON, *On a quadrature formula for trigonometric integrals*, Proc. Royal Soc. Edinburgh, 49 (1928), pp. 38–47.

Interpolate only the function $f(x)$ at $c_1 h, \dots, c_\nu h$ by a polynomial $\bar{f}(x)$

$$I[f] \approx Q_\nu^F[f] = \int_0^h \bar{f}(x) e^{i\omega x} dx = h \sum_{l=1}^{\nu} b_l(ih\omega) f(c_l h)$$

$$b_l(ih\omega) = \int_0^1 \ell_l(x) e^{ih\omega x} dx$$

ℓ_l is the l th cardinal polynomial of Lagrangian interpolation.

1-node Filon-type rule

$$I[f] = \int_0^h F(x) dx = \int_0^h f(x) \exp(i\omega x) dx$$

$$Q_1^F[f] = \frac{\exp(ih\omega) - 1}{i\omega} f(c_1 h)$$

$$Q_1^{EF}[F] = \frac{e^{ih\omega} - 1}{i\omega} f(h/2)$$

$$Q_1^F[f] = Q_1^{EF}[F] \text{ iff } c_1 = \frac{1}{2}$$

1-node Filon-type rule

$$I[f] = \int_0^h F(x) dx = \int_0^h f(x) \exp(i\omega x) dx$$

$$Q_1^F[f] = \frac{\exp(ih\omega) - 1}{i\omega} f(c_1 h)$$

$$Q_1^{EF}[F] = \frac{e^{ih\omega} - 1}{i\omega} f(h/2)$$

$$Q_1^F[f] = Q_1^{EF}[F] \text{ iff } c_1 = \frac{1}{2}$$

1-node Filon-type rule

$$I[f] = \int_0^h F(x) dx = \int_0^h f(x) \exp(i\omega x) dx$$

$$Q_1^F[f] = \frac{\exp(ih\omega) - 1}{i\omega} f(c_1 h)$$

$$Q_1^{EF}[F] = \frac{e^{ih\omega} - 1}{i\omega} f(h/2)$$

$$Q_1^F[f] = Q_1^{EF}[F] \text{ iff } c_1 = \frac{1}{2}$$

2-node Filon-type rule

$$I[f] = \int_0^h F(x) dx = \int_0^h f(x) \exp(i\omega x) dx$$

If f is interpolated at $c_1 h$ and $c_2 h$, then

$$\begin{aligned} Q_2^F[f] = & h \left[\left(\frac{i((e^{i\psi} - 1)c_2 - e^{i\psi})}{(c_1 - c_2)\psi} + \frac{e^{i\psi} - 1}{(c_1 - c_2)\psi^2} \right) f(c_1 h) \right. \\ & \left. + \left(\frac{i((e^{i\psi} - 1)c_1 - e^{i\psi})}{(c_2 - c_1)\psi} + \frac{e^{i\psi} - 1}{(c_2 - c_1)\psi^2} \right) f(c_2 h) \right] \end{aligned}$$

$Q_2^F[f] = Q_2^{EF}[F]$ iff the same nodes are used

2-node Filon-type rule

$$I[f] = \int_0^h F(x) dx = \int_0^h f(x) \exp(i\omega x) dx$$

If f is interpolated at $c_1 h$ and $c_2 h$, then

$$Q_2^F[f] = h \left[\left(\frac{i((e^{i\psi} - 1)c_2 - e^{i\psi})}{(c_1 - c_2)\psi} + \frac{e^{i\psi} - 1}{(c_1 - c_2)\psi^2} \right) f(c_1 h) \right. \\ \left. + \left(\frac{i((e^{i\psi} - 1)c_1 - e^{i\psi})}{(c_2 - c_1)\psi} + \frac{e^{i\psi} - 1}{(c_2 - c_1)\psi^2} \right) f(c_2 h) \right]$$

$Q_2^F[f] = Q_2^{EF}[F]$ iff the same nodes are used

Accuracy of Filon-type rules

A. Iserles, *On the numerical quadrature of highly-oscillating integrals. I. Fourier transforms*, IMA J. Numer. Anal., 24 (2004), pp. 365–391.

For small ω , a Filon-type quadrature method has an order as if $\omega = 0$.

Legendre nodes : order 2ν

Lobatto nodes : order $2\nu - 2$

For large ω :

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

Accuracy of Filon-type rules

A. Iserles, *On the numerical quadrature of highly-oscillating integrals. I. Fourier transforms*, IMA J. Numer. Anal., 24 (2004), pp. 365–391.

For small ω , a Filon-type quadrature method has an order as if $\omega = 0$.

Legendre nodes : order 2ν

Lobatto nodes : order $2\nu - 2$

For large ω :

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

Accuracy of Filon-type rules

A. Iserles, *On the numerical quadrature of highly-oscillating integrals. I. Fourier transforms*, IMA J. Numer. Anal., 24 (2004), pp. 365–391.

For small ω , a Filon-type quadrature method has an order as if $\omega = 0$.

Legendre nodes : order 2ν

Lobatto nodes : order $2\nu - 2$

For large ω :

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

Accuracy of Filon-type rules

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

$$\begin{aligned} Q_\nu^F[f] - I[f] &= I[\bar{f}] - I[f] = I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right] \end{aligned}$$

If $(c_1, c_\nu) = (0, 1)$ then $v(h) = v(0) = 0$

$$\implies Q_\nu^F[f] - I[f] = O(\omega^{-2}).$$

Accuracy of Filon-type rules

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

$$\begin{aligned} Q_\nu^F[f] - I[f] &= I[\bar{f}] - I[f] = I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right] \end{aligned}$$

If $(c_1, c_\nu) = (0, 1)$ then $v(h) = v(0) = 0$

$\implies Q_\nu^F[f] - I[f] = O(\omega^{-2})$.

Accuracy of Filon-type rules

$$Q_\nu^F[f] - I[f] \sim \begin{cases} O(\omega^{-1}) & c_1 > 0 \text{ or } c_\nu < 1 \\ O(\omega^{-2}) & c_1 = 0, c_\nu = 1 \end{cases}$$

$$\begin{aligned} Q_\nu^F[f] - I[f] &= I[\bar{f}] - I[f] = I[v] \\ &= - \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right] \end{aligned}$$

If $(c_1, c_\nu) = (0, 1)$ then $v(h) = v(0) = 0$

$$\implies Q_\nu^F[f] - I[f] = O(\omega^{-2}).$$

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p-1$$

$$Q_\nu^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p-1$$

$$Q_\nu^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p - 1$$

$$Q_{\nu}^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p - 1$$

$$Q_{\nu}^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p - 1$$

$$Q_{\nu}^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:

$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p - 1$$

$$Q_{\nu}^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

How to improve the accuracy of Filon-rules ?

- by using **Hermite interpolation** : asymptotic order $p + 1$ can be reached where p is the number of derivatives at the endpoints:



$$\bar{f}^{(l)}(h) = f^{(l)}(h), \bar{f}^{(l)}(0) = f^{(l)}(0), l = 0, \dots, p - 1$$

$$Q_{\nu}^F[f] - I[f] = O(\omega^{-p-1})$$

- by using **adaptive Filon-type methods** : allowing the interpolation points to depend on ω (is discussed later)
- by using **nodes in the complex plane** (=method of steepest descent)

Method of steepest descent

D. HUYBRECHS AND S. VANDEWALLE, *On the evaluation of highly oscillatory integrals by analytic continuation*, SIAM J. Numer. Anal., 44 (2007) pp 1026–1048.



Method of steepest descent

$$\begin{aligned} & \int_a^b f(x) e^{i\omega x} dx \\ &= e^{i\omega a} \int_0^\infty f(a + ip) e^{-\omega p} dp - e^{i\omega b} \int_0^\infty f(b + ip) e^{-\omega p} dp \\ &= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq \end{aligned}$$

This leads to the numerical evaluation of the two resulting integrals with classical **Gauss-Laguerre quadrature**.

High asymptotic order is obtained : using ν points for each integral, the error behaves as $O(\omega^{-2\nu-1})$.

Method of steepest descent

$$\begin{aligned} & \int_a^b f(x) e^{i\omega x} dx \\ &= e^{i\omega a} \int_0^\infty f(a + ip) e^{-\omega p} dp - e^{i\omega b} \int_0^\infty f(b + ip) e^{-\omega p} dp \\ &= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq \end{aligned}$$

This leads to the numerical evaluation of the two resulting integrals with classical **Gauss-Laguerre quadrature**.

High asymptotic order is obtained : using ν points for each integral, the error behaves as $O(\omega^{-2\nu-1})$.

Method of steepest descent

$$\begin{aligned}
 & \int_a^b f(x) e^{i\omega x} dx \\
 &= e^{i\omega a} \int_0^\infty f(a + ip) e^{-\omega p} dp - e^{i\omega b} \int_0^\infty f(b + ip) e^{-\omega p} dp \\
 &= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq
 \end{aligned}$$

This leads to the numerical evaluation of the two resulting integrals with classical **Gauss-Laguerre quadrature**.

High asymptotic order is obtained : using ν points for each integral, the error behaves as $O(\omega^{-2\nu-1})$.

Method of steepest descent

$$\begin{aligned}
 & \int_a^b f(x) e^{i\omega x} dx \\
 &= e^{i\omega a} \int_0^\infty f(a + ip) e^{-\omega p} dp - e^{i\omega b} \int_0^\infty f(b + ip) e^{-\omega p} dp \\
 &= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq
 \end{aligned}$$

This leads to the numerical evaluation of the two resulting integrals with classical **Gauss-Laguerre quadrature**.

High asymptotic order is obtained : using ν points for each integral, the error behaves as $O(\omega^{-2\nu-1})$.

Method of steepest descent

$$\int_a^b f(x) e^{i\omega x} dx$$

$$= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq$$

One ends up evaluating f at the points

$$a + i\frac{x_{nj}}{\omega}, \text{ and } b + i\frac{x_{nj}}{\omega}, \quad j = 1, \dots, n,$$

where x_{nj} are the n roots of the Laguerre polynomial of degree n .

This approach is equivalent to using a Filon rule with the same interpolation points.

Method of steepest descent

$$\int_a^b f(x) e^{i\omega x} dx$$

$$= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq$$

One ends up evaluating f at the points

$$a + i\frac{x_{nj}}{\omega}, \text{ and } b + i\frac{x_{nj}}{\omega}, \quad j = 1, \dots, n,$$

where x_{nj} are the n roots of the Laguerre polynomial of degree n .

This approach is equivalent to using a Filon rule with the same interpolation points.

Method of steepest descent

$$\int_a^b f(x) e^{i\omega x} dx$$

$$= \frac{e^{i\omega a}}{\omega} \int_0^\infty f\left(a + i\frac{q}{\omega}\right) e^{-q} dq - \frac{e^{i\omega b}}{\omega} \int_0^\infty f\left(b + i\frac{q}{\omega}\right) e^{-q} dq$$

One ends up evaluating f at the points

$$a + i\frac{x_{nj}}{\omega}, \text{ and } b + i\frac{x_{nj}}{\omega}, \quad j = 1, \dots, n,$$

where x_{nj} are the n roots of the Laguerre polynomial of degree n .

This approach is equivalent to using a Filon rule with the same interpolation points.

Adaptive Filon-type rules

Idea : combine best properties of EF and Filon quadrature

- EF
 - + accurate for small ωh since the method reduces to Gauss-Legendre quadrature
 - + good results for large ωh since the nodes tend to the endpoints (at a rate proportional to ω^{-1})
 - but : difficult to determine the nodes and weights for a given ωh (iteration needed and ill-conditioned)
- Filon
 - + any set of nodes can be used
 - there is no optimal set of nodes for all ωh
 - most accurate for small ωh if the method is built on Legendre nodes
 - most accurate for large ωh if the endpoints are included in the set of nodes

Adaptive Filon-type rules

Idea : combine best properties of EF and Filon quadrature

- EF
 - + accurate for small ωh since the method reduces to Gauss-Legendre quadrature
 - + good results for large ωh since the nodes tend to the endpoints (at a rate proportional to ω^{-1})
 - but : difficult to determine the nodes and weights for a given ωh (iteration needed and ill-conditioned)
- Filon
 - + any set of nodes can be used
 - there is no optimal set of nodes for all ωh
 - most accurate for small ωh if the method is built on Legendre nodes
 - most accurate for large ωh if the endpoints are included in the set of nodes

Adaptive Filon-type rules

Idea : combine best properties of EF and Filon quadrature

- EF
 - + accurate for small ωh since the method reduces to Gauss-Legendre quadrature
 - + good results for large ωh since the nodes tend to the endpoints (at a rate proportional to ω^{-1})
 - but : difficult to determine the nodes and weights for a given ωh (iteration needed and ill-conditioned)
- Filon
 - + any set of nodes can be used
 - there is no optimal set of nodes for all ωh
 - most accurate for small ωh if the method is built on Legendre nodes
 - most accurate for large ωh if the endpoints are included in the set of nodes

Adaptive Filon-type rules

Idea : combine best properties of EF and Filon quadrature

- EF
 - + accurate for small ωh since the method reduces to Gauss-Legendre quadrature
 - + good results for large ωh since the nodes tend to the endpoints (at a rate proportional to ω^{-1})
 - but : difficult to determine the nodes and weights for a given ωh (iteration needed and ill-conditioned)
- Filon
 - + any set of nodes can be used
 - there is no optimal set of nodes for all ωh
 - most accurate for small ωh if the method is built on Legendre nodes
 - most accurate for large ωh if the endpoints are included in the set of nodes

Adaptive Filon-type rules

Idea : create quadrature rules with ω -dependent nodes that

- reduce to Legendre-nodes for small ω
- reduce to Lobatto-nodes for large ω
- for given value of ω are easy to compute

To do so, we introduce S-shaped functions.

Adaptive Filon-type rules

Idea : create quadrature rules with ω -dependent nodes that

- reduce to Legendre-nodes for small ω
- reduce to Lobatto-nodes for large ω
- for given value of ω are easy to compute

To do so, we introduce S-shaped functions.

Adaptive Filon-type methods

- $\nu = 2$: $c_1(\psi) = \frac{3 - \sqrt{3}}{6} S(\psi; 2\pi; 1)$; $c_2(\psi) = 1 - c_1(\psi)$
- $\nu = 3$: $c_1(\psi) = \frac{10 - \sqrt{15}}{5} S(\psi; 3\pi; 1)$; $c_3(\psi) = 1 - c_1(\psi)$




Figure: $c_2(\psi)$ of the adaptive Filon method Q_2^{F-A} and $c_3(\psi)$ of the adaptive Filon method Q_3^{F-A} .

Asymptotic analysis for Q_2^{F-A}

$\tilde{c}_1 = c_1 h = \sigma_1(\omega)$ and $\tilde{c}_2 = c_2 h = h + \sigma_2(\omega)$ with $\sigma_{1,2}(\omega) \sim \omega^{-1}$

$$v(x) = s_h(x)(x - h - \sigma_2) \quad s_h(x) = \frac{f''(\xi_h(x))}{2}(x - \sigma_1)$$

$$v'(x) = s_h(x) + s'_h(x)(x - h - \sigma_2)$$

$$v''(x) = 2s'_h(x) + s''_h(x)(x - h - \sigma_2)$$

$$\vdots$$

$$v(h) = -s_h(h)\sigma_2$$

$$v'(h) = s_h(h) - s'_h(h)\sigma_2$$

$$v''(h) = 2s'_h(h) - s''_h(h)\sigma_2$$

$$\vdots$$

Similar results for the other endpoint.



Asymptotic analysis for Q_2^{F-A}

$\tilde{c}_1 = c_1 h = \sigma_1(\omega)$ and $\tilde{c}_2 = c_2 h = h + \sigma_2(\omega)$ with $\sigma_{1,2}(\omega) \sim \omega^{-1}$

$$v(x) = s_h(x)(x - h - \sigma_2) \quad s_h(x) = \frac{f''(\xi_h(x))}{2}(x - \sigma_1)$$

$$v'(x) = s_h(x) + s'_h(x)(x - h - \sigma_2)$$

$$v''(x) = 2s'_h(x) + s''_h(x)(x - h - \sigma_2)$$

$$\vdots$$

$$v(h) = -s_h(h)\sigma_2$$

$$v'(h) = s_h(h) - s'_h(h)\sigma_2$$

$$v''(h) = 2s'_h(h) - s''_h(h)\sigma_2$$

$$\vdots$$

Similar results for the other endpoint.

Asymptotic analysis for Q_2^{F-A}

$\tilde{c}_1 = c_1 h = \sigma_1(\omega)$ and $\tilde{c}_2 = c_2 h = h + \sigma_2(\omega)$ with $\sigma_{1,2}(\omega) \sim \omega^{-1}$

$$v(x) = s_h(x)(x - h - \sigma_2) \quad s_h(x) = \frac{f''(\xi_h(x))}{2}(x - \sigma_1)$$

$$v'(x) = s_h(x) + s'_h(x)(x - h - \sigma_2)$$

$$v''(x) = 2s'_h(x) + s''_h(x)(x - h - \sigma_2)$$

$$\vdots$$

$$v(h) = -s_h(h)\sigma_2$$

$$v'(h) = s_h(h) - s'_h(h)\sigma_2$$

$$v''(h) = 2s'_h(h) - s''_h(h)\sigma_2$$

$$\vdots$$

Similar results for the other endpoint.

Asymptotic analysis for Q_2^{F-A}

$$Q_2^{F-A}[f] - I[f] = I[v] \sim \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right]$$

Reordering for $s_h(h)$, $s'_h(h)$, ...

$$\begin{aligned} I[v] \sim & s_h(h) e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h) e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ & + s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

$$\sigma_2 = -\sigma_1 \text{ with } \sigma_{1,2}(\omega) \sim \psi^{-1} \iff Q_2^{F-A}[f] - I[f] \sim O(\psi^{-2})$$

Asymptotic analysis for Q_2^{F-A}

$$Q_2^{F-A}[f] - I[f] = I[v] \sim \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right]$$

Reordering for $s_h(h)$, $s'_h(h)$, ...

$$\begin{aligned} I[v] \sim & s_h(h) e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h) e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ & + s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

$$\sigma_2 = -\sigma_1 \text{ with } \sigma_{1,2}(\omega) \sim \psi^{-1} \iff Q_2^{F-A}[f] - I[f] \sim O(\psi^{-2})$$

Asymptotic analysis for Q_2^{F-A}

$$Q_2^{F-A}[f] - I[f] = I[v] \sim \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right]$$

Reordering for $s_h(h)$, $s'_h(h)$, ...

$$\begin{aligned} I[v] \sim & s_h(h) e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h) e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ & + s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

$$\sigma_2 = -\sigma_1 \text{ with } \sigma_{1,2}(\omega) \sim \psi^{-1} \iff Q_2^{F-A}[f] - I[f] \sim O(\psi^{-2})$$

Asymptotic analysis for Q_2^{F-A}

$$Q_2^{F-A}[f] - I[f] = I[v] \sim \sum_{m=0}^{\infty} \frac{1}{(-i\omega)^{m+1}} \left[e^{i\omega h} v^{(m)}(h) - v^{(m)}(0) \right]$$

Reordering for $s_h(h)$, $s'_h(h)$, ...

$$\begin{aligned} I[v] \sim & s_h(h) e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h) e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ & + s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

$$\sigma_2 = -\sigma_1 \text{ with } \sigma_{1,2}(\omega) \sim \psi^{-1} \iff Q_2^{F-A}[f] - I[f] \sim O(\psi^{-2})$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned} I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned}
 I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\
 &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots
 \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned}
 I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\
 &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots
 \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned} I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned} I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

A complex adaptive Filon-rule : Q_2^{F-C}

Are there better options than choosing $\sigma_2 = -\sigma_1$?

$$\begin{aligned} I[v] &\sim s_h(h)e^{i\psi} \left[\frac{\sigma_2}{i\omega} - \frac{1}{\omega^2} \right] + s'_h(h)e^{i\psi} \left[\frac{\sigma_2}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \\ &+ s_0(0) \left[\frac{\sigma_1}{i\omega} - \frac{1}{\omega^2} \right] + s'_0(0) \left[\frac{\sigma_1}{\omega^2} + \frac{2}{i\omega^3} \right] + \dots \end{aligned}$$

Yes : Suppose $\sigma_1 = \sigma_2 = i/\omega \implies Q_2^{F-C}[f] - I[f] \sim O(\psi^{-3})$.

$$Q_2^{F-C} = \frac{ih \left[f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi) \right]}{\psi}, \quad \psi = \omega h$$

Illustration




Figure: The normalised errors in some $\nu = 2$ Filon-type schemes for $f(x) = e^x$, $h = 1/10$ and different values of ω .

Error control for Q_2^{F-C}

$$Q_2^{F-C} = \frac{ih [f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi)]}{\psi}, \quad \psi = \omega h.$$

Obtained by replacing f by interpolating polynomial \bar{f} in nodes $i h/\omega$ and $h + i h/\omega$ (for large $\psi : \sim \psi^{-3}$)

Similarly : Q_3^{F-C} by replacing f by interpolating polynomial \tilde{f} in nodes $i h/\omega$, $h/2$ and $h + i h/\omega$ (for large ψ : also $\sim \psi^{-3}$ but about 100 times more accurate)

$$I[f] - I[\bar{f}] \approx I[\tilde{f}] - I[\bar{f}] = \frac{(1 - e^{i\psi})2h}{\psi^2(4 + \psi^2)} \times \\ \left((2 - i\psi) f\left(\frac{i}{\omega}\right) - (2 + i\psi) f\left(h + \frac{i}{\omega}\right) + (2i\psi) f\left(\frac{h}{2}\right) \right)$$

Error control for Q_2^{F-C}

$$Q_2^{F-C} = \frac{ih [f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi)]}{\psi}, \quad \psi = \omega h.$$

Obtained by replacing f by interpolating polynomial \bar{f} in nodes $i h/\omega$ and $h + i h/\omega$ (for large $\psi : \sim \psi^{-3}$)

Similarly : Q_3^{F-C} by replacing f by interpolating polynomial \tilde{f} in nodes $i h/\omega$, $h/2$ and $h + i h/\omega$ (for large ψ : also $\sim \psi^{-3}$ but about 100 times more accurate)

$$I[f] - I[\bar{f}] \approx I[\tilde{f}] - I[\bar{f}] = \frac{(1 - e^{i\psi})2h}{\psi^2(4 + \psi^2)} \times \\ \left((2 - i\psi) f\left(\frac{i}{\omega}\right) - (2 + i\psi) f\left(h + \frac{i}{\omega}\right) + (2i\psi) f\left(\frac{h}{2}\right) \right)$$

Error control for Q_2^{F-C}

$$Q_2^{F-C} = \frac{ih [f(ih/\psi) - e^{i\psi} f((i+\psi)h/\psi)]}{\psi}, \quad \psi = \omega h.$$

Obtained by replacing f by interpolating polynomial \bar{f} in nodes $i h/\omega$ and $h + i h/\omega$ (for large $\psi : \sim \psi^{-3}$)

Similarly : Q_3^{F-C} by replacing f by interpolating polynomial \tilde{f} in nodes $i h/\omega$, $h/2$ and $h + i h/\omega$ (for large ψ : also $\sim \psi^{-3}$ but about 100 times more accurate)

$$I[f] - I[\bar{f}] \approx I[\tilde{f}] - I[\bar{f}] = \frac{(1 - e^{i\psi})2h}{\psi^2(4 + \psi^2)} \times \\ \left((2 - i\psi) f\left(\frac{i}{\omega}\right) - (2 + i\psi) f\left(h + \frac{i}{\omega}\right) + (2i\psi) f\left(\frac{h}{2}\right) \right)$$



Illustration




Figure: Error estimations for the Q_2^{F-A} and Q_2^{F-C} method applied on the problem with $f(x) = e^x$, $h = 2$.

Illustration




Figure: Error estimations for the Q_2^{F-A} and Q_2^{F-C} method applied on the problem with $f(x) = e^x$, $h = 2$.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.

Conclusions

- Filon rules, EF rules, and steepest descent rules are built up starting from different points of view, the basic underlying idea is the same : $f(x)$ is interpolated by a polynomial.
- Different choices can be made for the interpolation nodes.
- A choice of the (complex) interpolation nodes can improve the asymptotic behaviour of the quadrature rule.
- Even better asymptotic behaviour is obtained if the nodes are frequency dependent.
- Cheap error estimation is possible.